

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Patent Application of:
Marche and al.

Confirmation No.: 7396

Serial No.: 10/581,814

Art Unit: 1637

Filed: December 3, 2004

Examiner: Samuel Woolwine

For: METHOD FOR QUANTITATIVE EVALUATION OF A REARRANGEMENT OR A
TARGETED GENETIC RECOMBINATION OF AN INDIVIDUAL AND USES
THEREOF

DECLARATION UNDER RULE 132

Commissioner of Patents
PO BOX 1450
Alexandria, VA 22313-1450

Dear Commissioner:

Now comes Nicolas Pasqual, who declares and states that:

1. I have the following degrees and qualifications: PhD in Biology and immunology, M.S. in Cellular & Molecular Biology, and a B.S. in physiology.
2. I have been employed by COMMISSARIAT A L'ENERGIE ATOMIQUE ET AUX ENERGIES ALTERNATIVES (CEA) for 3 years and I am now President, CEO and CSO of ImmunID Technologies (since 5 years).
3. I declare that I am experienced in the field of Immunology, immunogenetic, molecular biology and biology, as may be seen from the attached truncated curriculum vitae and the publication list attached hereto.
4. I am one of the named inventor of the above-identified Patent Application ("the '814 Application")
5. I have reviewed the Office Action dated October 30, 2009 and I declare the following :
6. Contrary to what was described in Pasqual and al. the instant method by combining the followings steps of:

- Amplifying a **long segment of gDNA** (having a size between a few hundred base pairs and several tens of kb in size) from a **human biological sample** by long multiplex PCR in which the elongation steps are performing at least for 10 minutes at 68°-72°C,
- Using at least one pair of primers in which (1) one of said primers V hybridizes upstream or at the 5'terminal end of a V gene of the variable domain of the α chain of TCR and (ii) the other primer J hybridizes downstream or at the 3' end of a gene J of the α chain of TCR,
- Using a DNA polymerase having a correction activity, and
- Separating the amplified DNA products by electrophoretic migration on a gel and detecting said amplified products directly on said gel by using a DNA-labeling agent during the migration and detecting after excitation in the UV range or another appropriate wave length or by using during amplification, labeled primers,

is able unexpectedly to improve the readability of the gels compared to the results observed in Pasqual and al. conditions, in which a Southern blot is compulsory in view to obtain readable results.

7. Tests were performed under my supervision and confirm that it is only by combining the specific features of the instant invention that it is possible to avoid the use of the Southern blot step which is not adapted to routine tests.

8. Exhibit A : Pasqual et al. conditions

In the conditions of Pasqual et al., it was possible, at the time of the publication of Pasqual et al., to observe and resolve 4 to 5 rearrangements by multiplex PCR, only when the DNA was transferred on a membrane and radioactive probes were used (figure 2A), whereas in figure 1A, which corresponds to results obtained directly on a gel, only smears (which cannot be interpreted) are observed.

In these two figures (1A and 2A) 6 TCRAD V genes have been studied (V2, V12, V19, V20 and V101).

The results observed in figure 1A are not due to a coloration problem since the size markers (right lane of the gel in figure 1A) are clearly visible.

It emerges from said figures that in the conditions of Pasqual et al, it is not possible to measure the combinatorial diversity of the immune repertoire TRAV-J, directly on a gel using a fluorescent detection. In such a case, the very weak "specific signal" is hidden in a smear of non specific signal also very weak (see figure 1A). Therefore it is not possible to detect a V-J rearrangement directly after PCR. The only possibility, in the conditions of Pasqual et al. , to obtain a detectable signal, was to perform a Southern blot using radioactive probes (see figure 2A).

On figure 2A, the black blots correspond to the rearrangements detected by the radioactive probes used in the Southern Blot.

Such a method is very long to carry out, costly and risky:

- . Transfer of the amplicons to a membrane (>4h)
- . Fixation of the DNA on the membrane (some minutes)
- . Labeling of the probes (20-60 minutes)

- . Incubation in a hybridization roller (on the membrane (> 2h)
- . Washings (30-120 minutes)
- . Exposure of the membrane in a cassette in view to retrieve the radioactive signal (> 4H-several days)
- . Reading the cassette (15 minutes).

It is not adapted to routine diagnostic tests.

9. Exhibit B : Pasqual et al. conditions

Method of Multiplex PCR assay:

Briefly, using an upstream primer specific for a given Vx family and a downstream primer specific for a given Jy segment, the multiplex PCR assay allows the detection of a Vx to Jy rearrangement as well as that of Vx genes to a limited set of 5' J segments spanning from Jy to Jy-4 position.

Amplifications were performed with 1.3 unit/reaction of Expend High Fidelity PCR system (Roche Diagnostics, Meylan, France)

The cycling conditions were 5 min at 94°C, 26 cycles of 1 min at 94°C, 1 min at 58°C, 6 min at 72°C, and one cycle of 10 min at 72°C. In these assay conditions, maximum amplicon size was approximately 5kb.

Quality control PCR using a non rearranged gene P53.

Normalization of the quantity of DNA in each reaction was determined by amplification of p53 gene **as a non-rearranging gene** in the same PCR run.

PCR is done using P53 specific primers. Primer P53 sens X6.5 : ACAGCGTGGTGGTACCTTAT, and Primer P53 anti-sens X7 : CACATGTACTTGTAAGTGGATGG.

Migration : 10 µL of amplicon + 2 µL of loading dye is loaded on agarose Gel (1,5%) + ethidium bromide. Migration at 150V during 3hours. UV excitation in a Vilber Lourmat + camera box.

Results on Gel

P53 normalization: figure 3B

Direct detection of P53 gene (after UV excitation) in each sample indicates a good normalization of DNA quantity for each sample F18, F19, F20, J0, J1 and J28.

This information confirms that gDNA quantity and quality is good, thus rearrangement detection can be performed on these samples.

Secondarily this result indicates that Gel coloration is good and if no rearrangement can be directly detected on gel it is not a matter of gel coloration and UV excitation.

Rearrangements: figures 1B and 2B

Smear detection (after UV excitation) and quasi non distinguished V-J rearrangements **on Gel** for both ADV19 (figure 2B) and ADV20 genes (figure 3B) on J56, J48, J40, J33, J27, J16, J9 and J2.

It is compulsory to perform an additional Southern Blot with internal radioactive probe (figures 4B and 5B).

Results are interpretable only on Southern blotted membrane (figures 4B and 5B)

Detection of ADV19 rearrangements on central and distal J genes at J1 and J28: figure 4B

Detection of ADV20 rearrangement on proximal J genes at F19, F20, J0, J1, and J28 figure 5B.

Conclusions

The results observed are not due to coloration problem since the size markers (right lane of the gel in figures 1B and 2B) are clearly visible.

Furthermore whereas in figures 1B and 2B smears are obtained, in figure 3B, good results are obtained directly on a gel, when in the same conditions, as in figures 1B and 2B, gene P53 is used.

Thus in the conditions of document Pasqual and al., a step of Southern blot is compulsory in view to characterize the rearrangements.

On the contrary, in the conditions of the instant invention, it is possible to obtain a map of the immune repertoire with strong signals, this allowing a direct detection on gels with the possibility of using such a method in routine diagnostic tests, in less than one day.

10. Exhibit C: Instant invention conditions

Figure 1C illustrates the results obtained from a human sample, with the method according to the instant invention.

More precisely:

Multiplex PCR assay: multiplex PCR is performed with V1 (=ADV19) and V8 (ADV18) upstream primers on J5 and J33 gene downstream primers as described in the present invention.

Migration : 10 μ L of amplicon + 2 μ L of loading dye is loaded on agarose Gel (1,5%) + ethidium bromide. Migration at 150V during 3hours. UV excitation in a Vilber Lourmat + camera box.

Results : Direct UV detection of V1 and V8 rearrangements on J5 and upstream J genes as well as J33 counterparts.

Coloration with SyberGreen (or BET) is possible and effectively allows the detection of the rearrangements directly on the gel.

11. I declare that all statements made herein of my own knowledge are true and that all statements made on information and belief to be true; and that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the Application or an Patent issuing there from.

Date: 12/4/2010



Nicolas PASQUAL

Exhibit A




Figure 1




Exhibit B

Page

Figure 2



Figure 4

Postcard

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

526

527

528

529

530

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

ADV 20-74T

Figure 5

Figure 3

Exhibit C




Figure 1